

Appendix 2A

Study	Year	Lesion Location	Number of Lesions	Average Size of Lesion (SD/Range)
Karthikeyan <i>et al.</i> ²⁰	2012	Ac (Sup and Ant-Sup zones)	1	154mm ² (50-300mm ²)
McDonald <i>et al.</i> ²¹	2014	Ac or FH (Outerbridge grade IV)	NR	119mm ² (20-250mm ²)
McDonald <i>et al.</i> ²²	2013	30 Ac, 5 FH, 4 Ac+FH	NR	162mm ² (20-378mm ²)
Philippon <i>et al.</i> ²³	2008	Ac (Sup quadrant 9-3 o'clock) + FH in 1 Pt	1-2	163mm ² (40-240mm ²)
Singh <i>et al.</i> ²⁴	2010	22 cam, 3 pincer, 2 pincer+cam, 93% rim	1	“Lesions up to 300mm ² ”, mean NR

Appendix 2B

Study	Year	Complications	Reoperations
Karthikeyan <i>et al.</i> ²⁰	2012	1 failure: avascular necrosis	All cases were re-operation after an initial arthroscopy.
McDonald <i>et al.</i> ²¹	2014	NR	NR
McDonald <i>et al.</i> ²²	2013	NR	NR
Philippon <i>et al.</i> ²³	2008	12 failures: 6. capsulolabral adhesions; 3. myositis ossificans; 2. femoral head defect	All cases were re-operation after an initial arthroscopy. One patient underwent a total hip arthroplasty 3 years following secondary evaluation
Singh <i>et al.</i> ²⁴	2010	NR	NR